

## Brief review of oscillations

Assume a  $2 \times 2$  neutrino mixing matrix.

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}$$

$$\nu_a(t) = \cos(\theta)\nu_1(t) + \sin(\theta)\nu_2(t)$$

$$\begin{aligned} P(\nu_a \rightarrow \nu_b) &= |<\nu_b|\nu_a(t)>|^2 \\ &= \sin^2(\theta)\cos^2(\theta)|e^{-iE_2 t} - e^{-iE_1 t}|^2 \end{aligned}$$

Sufficient to understand most of the physics:

$$P(\nu_a \rightarrow \nu_b) = \sin^2 2\theta \sin^2 \frac{1.27((m_2^2 - m_1^2)/eV^2)(L/km)}{(E/GeV)}$$

**Appearance**

$$P(\nu_a \rightarrow \nu_a) = 1 - \sin^2 2\theta \sin^2 \frac{1.27(\Delta m^2/eV^2)(L/km)}{(E/GeV)}$$

**Disappearance**

Oscillation nodes at  $\pi/2, 3\pi/2, 5\pi/2, \dots (\pi/2)$ :  $\Delta m^2 = 0.0025eV^2$ ,  
 $E = 1GeV$ ,  $L = 494km$ .